

■文献リスト

ワサビ

- Shin IS, Masuda H, Naohide K. (2004) Bactericidal activity of wasabi (*Wasabia japonica*) against *Helicobacter pylori*. *Int J Food Microbiol.* ;94:255-61.
- Hasegawa N, Matsumoto Y, Hoshino A, Iwashita K. (1999) Comparison of effects of *Wasabia japonica* and allyl isothiocyanate on the growth of four strains of *Vibrio parahaemolyticus* in lean and fatty tuna meat suspensions. *Int J Food Microbiol.*;49:27-34.
- Yano T, Yajima S, Virgona N, Yano Y, Otani S, Kumagai H, Sakurai H, Kishimoto M, Ichikawa T. (2000) The effect of 6-methylthiohexyl isothiocyanate isolated from *Wasabia japonica* (wasabi) on 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol-induced lung tumorigenesis in mice. *Cancer Lett.*;155 :115-20.
- Makoto Watanabe, Masahiko Ohata, Sumio Hayakawa, Mamoru Isemura, Shigenori Kumazawa, Tsutomu Nakayama, Michiyo Furugori and Naohide Kinae (2003) Identification of 6-methylsulfinylhexyl isothiocyanate as an apoptosis-inducing component in wasabi. *Phytochemistry* ; 62 : 733-9.
- Morimitsu Y, Hayashi K, Nakagawa Y, Horio F, Uchida K, Osawa T. (2000) Antiplatelet and anticancer isothiocyanates in Japanese domestic horseradish, wasabi. *Biofactors*; 13:271-6.
- Morimitsu Y, Nakagawa Y, Hayashi K, Fujii H, Kumagai T, Nakamura Y, Osawa T, Horio F, Itoh K, Iida K, Yamamoto M, Uchida K. (2002) A sulforaphane analogue that potently activates the Nrf2-dependent detoxification pathway. *J Biol Chem.*;277: 3456-63.

ウド

- Sakamura S. and Kawano K. (1970)Cyanidin 3-xylosylgalactoside, an anthocyanin from *Aralia elata* and *Aralia cordata*. *Phytochemistry*. 9(5) 1147.
- Asada Y., Sakamoto K. and Furuya Y.(1994)A minor anthocyanin from cultured cells of *Aralia cordata*. *Phytochemistry*. 35 (6), 1471-1473.
- 村山徹、箭田浩士 (2002) うどのか酸化成分の同定とその含量の部位及び栽培法による差異. 平成13年度農業技術研究機構研究成果情報、東北農業研究成果情報
- Okuyama E, Nishimura S, Yamazaki M.(1991)Analgesic principles from *Aralia cordata* Thunb.*Chem Pharm Bull (Tokyo)*. .39(2):405-7.
- Okuyama E, Nishimura S, Ohmori S, Ozaki Y, Satake M, Yamazaki M. (1993)Analgesic component of *Notopterygium incisum* Ting.*Chem Pharm Bull (Tokyo)*. 41(5):926-9.

フキとふきのとう

- Matsuura H, Amano M, Kawabata J, Mizutani(2002) Isolation and Measurement of Quercetin Glucosides in Flower Buds of Japanese Butterbur (*Petasites japonicus* subsp. *gigantea* Kitam.). *Biosci Biotechnol Biochem.*,66(7):1571-5.
- 田崎 弘之、吉原 照彦、岩本 嗣 (2002) フキ酸の生物有機化学的解析とその利用に関する研究、(財) 北海道科学・産業振興財団研究成果報告書
- Hasa Y, Tazaki H. Related Articles(2004) Biosynthesis of Fukinolic Acid Isolated from *Petasites japonicus*. *Biosci Biotechnol Biochem.*, 68:2212-4.
- Iriye R, Furukawa K, Nishida R, Kim C, Fukami H.(1992) Isolation and synthesis of a new bio-antimutagen, petasiphenol, from scapes of *Petasites japonicum*. *Biosci Biotechnol Biochem.*, 56(11):1773-5.
- Mizushina Y, Kamisuki S, Kasai N, Ishidoh T, Shimazaki N, Takemura M, Asahara H, Linn S, Yoshida S, Koiwai O, Sugawara F, Yoshida H, Sakaguchi K. (2002)Pet-asiphenol: a DNA polymerase lambda inhibitor. *Biochemistry* . , 41(49):14463-71.
- Mizushina Y, Hirota M, Murakami C, Ishidoh T, Kamisuki S, Shimazaki N, Takemura M, Perpelescu M, Suzuki M, Yoshida H, Sugawara F, Koiwai O, Sakaguchi K.(2003) Some anti-chronic inflammatory compounds are DNA polymerase lambda-specific inhibitors. *Biochem Pharmacol.*, 66(10):1935-44.
- Mizushina Y, Ishidoh T, Kamisuki S, Nakazawa S, Takemura M, Sugawara F, Yoshida H, Sakaguchi K. (2003) Flavonoid glycoside: a new inhibitor of eukaryotic DNA polymerase alpha and a new carrier for inhibitor-affinity chromatography. *Biochem Biophys Res Commun.*, 301(2):480-7.
- Matsubara K, Mori M, Mizushina Y. (2004)Petasiphenol which inhibits DNA polymerase lambda activity is an inhibitor of in vitro angiogenesis. *Oncol Rep.*, 11(2):447-51.
- Niwa H, Ishiwata H, Yamada K. (1983) Separation and determination of macrocyclic pyrrolizidine alkaloids of the otonecine type present in the edible plant *Petasites japonicus* by reversed-phase high-performance liquid chromatography. *J Chromatogr.*, 257(1):146-50.
- Tobinaga S, Takeuchi N, Kasama T, Yamashita J, Aida Y, Kaneko (1983)Anti-histaminic and anti-allergic principles of *Petasites japonicus* Maxim. *Chem Pharm Bull (Tokyo)*., 31(2):745-8.

タラ

- Yoshikawa M, Harada E, Matsuda H, Murakami T, Yamahara J, Murakami N. (1993)Elatosides A and B, potent inhibitors of ethanol absorption in rats from the bark of *Aralia elata* Seem: the structure-activity relationships of oleanolic acid oligoglycosides.*Chem Pharm Bull (Tokyo)*.41(11):2069-71.
- Sakai S, Katsumata M, Satoh Y, Nagasao M, Miyakoshi M, Ida Y, Shoji J. (1994)Oleanolic acid saponins from root bark of *Aralia elata*.*Phytochemistry* .,35(5):1319-24.
- Satoh Y, Sakai S, Katsumata M, Nagasao M, Miyakoshi M, Ida Y, Shoji J. (1994)Oleanolic acid saponins from root-bark of *Aralia elata*.*Phytochemistry* .,36(1):147-52.
- Yoshikawa M, Matsuda H, Harada E, Murakami T, Wariishi N, Yamahara J, Murakami N. (1994)Elatoside E, a new hypoglycemic principle from the root cortex of *Aralia elata* Seem.: structure-related hypoglycemic activity of oleanolic acid glycosides. *Chem Pharm Bull (Tokyo)* .,42(6):1354-6.
- Yoshikawa M, Yoshizumi S, Ueno T, Matsuda H, Murakami T, Yamahara J, Murakami N.(1995) Medicinal foodstuffs. I. Hypoglycemic constituents from a garnish foodstuff "taranome," the young shoot of *Aralia elata* SEEM.: elatosides G, H, I, J, and K. *Chem Pharm Bull (Tokyo)* .,43(11):1878-82.
- Yoshikawa M, Murakami T, Harada E, Murakami N, Yamahara J, Matsuda H. (1996)Bioactive saponins and glycosides. VI. Elatosides A and B, potent inhibitors of ethanol absorption, from the bark of *Aralia elata* SEEM. (Araliaceae): the structure-requirement in oleanolic acid glucuronide-saponins for the inhibitory activity. *Chem Pharm Bull (Tokyo)* .,44(10):1915-22.
- Yoshikawa M, Murakami T, Harada E, Murakami N, Yamahara J, Matsuda H. (1996)Bioactive saponins and glycosides. VII. On the hypoglycemic principles from the root cortex of *Aralia elata* Seem.: structure related hypoglycemic activity of oleanolic acid oligoglycoside. *Chem Pharm Bull (Tokyo)* .,44(10):1923-7.
- Chung CK, Jung ME. (2003)Ethanol fraction of *Aralia elata* Seemann enhances antioxidant activity and lowers serum lipids in rats when administered with benzo(a)pyrene.*Biol Pharm Bull.* ,26(10):1502-4.
- Wang Z, Song S, Lu H, Chen G, Xu S, Sagara Y, Kitaoka N, Manabe M, Kodama H. (2003)Effect of three triterpenoid compounds isolated from root bark of *Aralia elata* on stimulus-induced superoxide generation and tyrosyl phosphorylation and translocation of p47(phox) and p67(phox) to cell membrane in human neutrophil.*Clin Chim Acta* .,336(1-2):65-72.

Tomatsu M, Ohnishi-Kameyama M, Shibamoto N.(2003)Aralin, a new cytotoxic protein from *Aralia elata*, inducing apoptosis in human cancer cells.Cancer Lett.,199(1):19-25.

Tomatsu M, Mujin T, Shibamoto N, Tashiro F, Ikuta A. (2004) Production of aralin, a selective cytotoxic lectin against human transformed cells, in callus culture of *Aralia elata*.*Planta Med.* ,70(5):469-71.

Tomatsu M, Kondo T, Yoshikawa T, Komono T, Adachi N, Kawasaki Y, Ikuta A, Tashiro F. (2004)An apoptotic inducer, aralin, is a novel type II ribosome-inactivating protein from *Aralia elata*.*Biol Chem.* ,385(9):819-27.

Saito S, Ebashi J, Sumita S, Furumoto T, Nagamura Y, Nishida K, Isiguro I.(1993)Comparison of cytoprotective effects of saponins isolated from leaves of *Aralia elata* Seem. (Araliaceae) with synthesized bisdesmosides of oleanolic acid and hederagenin on carbon tetrachloride-induced hepatic injury.*Chem Pharm Bull (Tokyo)*.,41(8):1395-401.

アシタバ

Kimura Y, Taniguchi M, Baba K. (2004)Antitumor and antimetastatic activities of 4-hydroxyderricin isolated from Angelica keiskei roots.*Planta Med.*,70(3):211-9.

Akihisa T, Tokuda H, Ukiya M, Iizuka M, Schneider S, Ogasawara K, Mukainaka T, Iwatsuki K, Suzuki T, Nishino H. (2003)Chalcones, coumarins, and flavanones from the exudate of Angelica keiskei and their chemopreventive effects.*Cancer Lett.*, 201(2):133-7.

Kimura Y, Baba K. (2003)Antitumor and antimetastatic activities of Angelica keiskei roots, part 1: Isolation of an active substance, xanthoangelol.*Int J Cancer.*,106:429-37.

Ogawa H, Nakashima S, Baba K. (2003)Effects of dietary Angelica keiskei on lipid metabolism in stroke-prone spontaneously hypertensive rats.*Clin Exp Pharmacol Physiol.*,30(4):284-8.

Park JC, Park JG, Kim HJ, Hur JM, Lee JH, Sung NJ, Chung SK, Choi JW. (2002) Effects of extract from Angelica keiskei and its component, cynaroside, on the hepatic bromobenzene-metabolizing enzyme system in rats.*Phytother Res.*,16:24-7.

Shimizu E, Hayashi A, Takahashi R, Aoyagi Y, Murakami T, Kimoto K.(1999) Effects of angiotensin I-converting enzyme inhibitor from Ashitaba (*Angelica keiskei*) on blood pressure of spontaneously hypertensive rats.*J Nutr Sci Vitaminol (Tokyo)*.,45(3):375-83.

Inamori Y, Baba K, Tsujibo H, Taniguchi M, Nakata K, Kozawa M.(1991) Antibacterial activity of two chalcones, xanthoangelol and 4-hydroxyderricin, isolated from the root of Angelica keiskei KOIDZUMI.*Chem Pharm Bull (Tokyo)*.39(6):1604-5.

Okuyama T, Takata M, Takayasu J, Hasegawa T, Tokuda H, Nishino A, Nishino H, Iwashima A.(1991) Anti-tumor-promotion by principles obtained from Angelica keiskei.*Planta Med.*,57(3):242-6.

Okuyama T, Takata M, Nishino H, Nishino A, Takayasu J, Iwashima A. (1990) Studies on the antitumor-promoting activity of naturally occurring substances. II. Inhibition of tumor-promoter-enhanced phospholipid metabolism by umbelliferous materials.*Chem Pharm Bull (Tokyo)*.,38(4):1084-6.

<http://www.takara-bio.co.jp/>

ワラビ、ゼンマイ

林美央、三輪章志 (2002) 能登の山菜を活用した特産品の開発 II 山菜の機能性の検索、石川県農業総合研究センター報告、 24:61 ~ 64

Potter DM, Baird MS. (2000) Carcinogenic effects of ptaquiloside in bracken fern and related compounds. *Br J Cancer*,83(7):914-20.

クサソテツ

木村俊之、鈴木雅博、山岸賢治、新本洋士 (2002) コゴミの活性酸素消去成分、平成13年度農業技術研究機構研究成果情報、東北農業研究成果情報

Kimura T, Suzuki M, Takenaka M, Yamagishi K, Shimamoto H. (2004)L-O-Caffeoylhomoserine from *Matteuccia struthiopteris*. *Phytochemistry*, 65(4):423-6.

コシアブラ

島根県農業試験場 (2003) 薬草などの栽培技術の確立と利用技術の開発、平成12~14年度重点的科学技術開発事業、研究成果報告書

東北農業研究センター畠地利用部 (2003) コシアブラの抗酸化成分と促成栽培法、2003年版研究案内最近の研究トピックス

ウコギ

Yamada N., Ogata T. and Onodera J. (1997) Antioxidant activity of *Acathopanax sieboldianum* Makino (Ukogi in Japanese). Proceedings of the 2nd International Conference on Bioradicals. (Edited by H.Kamada and H.Ohya), 169-171.

田渕三保子、田村朝子、山田則子 (2003) ウコギ (*Acathopanax sieboldianum* m) 葉投与が新生児期のストレプトゾトシン投与による2型糖尿病ラットの耐糖能に及ぼす影響、日本栄養・食糧学会誌、56 (4), 159-162.

Fujikawa T, Yamaguchi A, Morita I, Takeda H, Nishibe S.(1996)Protective effects of *Acanthopanax senticosus* Harms from Hokkaido and its components on gastric ulcer in restrained cold water stressed rats. *Biol Pharm Bull.*, 19(9),1227-30.

セリ

Zhang H, Piao H, Li M, Wu Z, Piao S, Jin Z. Related(1995) Protective effects of methanolic extract of *Oenanthe stolonifera* (Roxb.) Wall. ex DC. on myocardial injury induced by ischemia and reperfusion in rats. *Zhongguo Zhong Yao Za Zhi*., 20(1):44-6, 63.

Hasegawaa H., Fukasawa-Akadaa T., Okunob T., Niizekic M., Suzukia M. (2001) Anthocyanin accumulation and related gene expression in Japanese parsley (*Oenanthe stolonifera*, DC.) induced by low temperature. *J. of Plant Physiol.*, 158(1): 71-78.

ミツバ

Nakano Y, Matsunaga H, Saita T, Mori M, Katano M, Okabe H.(1998) Antiproliferative constituents in Umbelliferae plants II. Screening for polyacetylenes in some Umbelliferae plants, and isolation of panaxynol and falcarindiol from the root of *Heracleum moellendorffii*. *Biol Pharm Bull.* , 21(3):257-61.

Ha Won Kima, Akira Murakamib, Yoshimasa Nakamura and Hajime Ohigashi(2002) Screening of edible Japanese plants for suppressive effects on phorbol ester-induced superoxide generation in differentiated HL-60 cells and AS52 cells. *Cancer Letters* , 176(1):7-16.

アサツキ ノビル

- Ghahfarokhi MS, Goodarzi M, Abyaneh MR, Al-Tiraihi T, Seyedipour G. (2004) Morphological evidences for onion-induced growth inhibition of *Trichophyton rubrum* and *Trichophyton mentagrophytes*. *Fitoterapia*, 75,645-55.
- Singh A, Arora A, Shukla Y. (2004) Modulation of altered hepatic foci induction by diallyl sulphide in Wistar rats. *Eur J Cancer Prev*, 13(4):263-9.
- Arora A, Siddiqui IA, Shukla Y.(2004)Modulation of p53 in 7,12-dimethylbenz[a]anthracene-induced skin tumors by diallyl sulfide in Swiss albino mice.*Mol Cancer Ther*,3(11):1459-66.
- Babu PS, Srinivasan K. (1999) Renal lesions in streptozotocin-induced diabetic rats maintained on onion and capsaicin containing diets.*Nutr Biochem*, 10(8):477-83.
- Turner B, Molgaard C, Marckmann P. (2004) Effect of garlic (*Allium sativum*) powder tablets on serum lipids, blood pressure and arterial stiffness in normo-lipidemic volunteers: a randomised, double-blind, placebo-controlled trial.*Br J Nutr*,92(4):701-6.
- Yang J, Meyers KJ, van der Heide J, Liu RH. (2004) Varietal differences in phenolic content and antioxidant and antiproliferative activities of onions.*J Agric Food Chem*, 52(22):6787-93.
- Donaldson MS. (2004) Nutrition and cancer: A review of the evidence for an anti-cancer diet. *Nutr J*, 3(1):19.
- Herman-Antosiewicz A, Singh SV. (2004) Signal transduction pathways leading to cell cycle arrest and apoptosis induction in cancer cells by Allium vegetable-derived organosulfur compounds: a review.*Mutat Res*, 555(1-2):121-31.
- Arivazhagan S, Velmurugan B, Bhuvaneswari V, Nagini S. (2004) Effects of aqueous extracts of garlic (*Allium sativum*) and neem (*Azadirachta indica*) leaf on hepatic and blood oxidant-antioxidant status during experimental gastric carcinogenesis.*J Med Food*, 7(3):334-9.
- Perez-Severiano F, Rodriguez-Perez M, Pedraza-Chaverri J, Maldonado PD, Medina-Campos ON, Ortiz-Plata A, Sanchez-Garcia A, Villeda-Hernandez J, Galvan-Arzate S, Aguilera P, Santamaria A. (2004) S-Allylcysteine, a garlic-derived antioxidant, ameliorates quinolinic acid-induced neurotoxicity and oxidative damage in rats.*Neurochem Int*, 45(8):1175-83.
- Kiesewetter H, Jung F, Mrowietz C, Pindur G, Heiden M, Wenzel E, Gu LD. (1990) Effects of garlic on blood fluidity and fibrinolytic activity: a randomised, placebo-controlled, double-blind study. *Br J Clin Pract Suppl*, 69:24-9.
- Keiss HP, Dirsch VM, Hartung T, Hauffner T, Trueman L, Auger J, Kahane R, Vollmar AM.(2003) Garlic (*Allium sativum* L.) modulates cytokine expression in lipopolysaccharide-activated human blood thereby inhibiting NF-kappaB activity.*J Nutr*, 133(7):2171-5.
- Fleischauer AT, Poole C, Arab L. (2000) Garlic consumption and cancer prevention: meta-analyses of colorectal and stomach cancers. *Am J Clin Nutr*, 72(4):1047-52.

ギョウジャニンニク

- Nishimura H, Takahashi T, Wijaya CH, Satoh A, Ariga T.(2000) Thermochemical transformation of sulfur compounds in Japanese domestic Allium, *Allium victorialis* L.*Biofactors*,13:257-63
- Shirataki Y, Motohashi N, Tani S, Sunaga K, Sakagami H, Satoh K, Nakashima H, Kanamoto T, Wolfard K, Molnar J. (2001) Antioxidative activity of *Allium victorialis* L. extracts. *Anticancer Res*,21:3331-9
- Lee KT, Choi JH, Kim DH, Son KH, Kim WB, Kwon SH, Park HJ. (2001) Constituents and the antitumor principle of *Allium victorialis* var. *platyphyllum*. *Arch Pharm Res*,24:44-50.

オオバギボウシ

- Mimaki Y, Kuroda M, Kameyama A, Yokosuka A, Sashida Y. (1998) Steroidal saponins from the rhizomes of *Hosta sieboldii* and their cytostatic activity on HL-60 cells.*Phytochemistry*,48(8):1361-9.
- Mimaki Y, Kameyama A, Kuroda M, Sashida Y, Hirano T, Oka K, Koike K, Nikaido T (1997) Steroidal glycosides from the underground parts of *Hosta plantaginea* var. *japonica* and their cytostatic activity on leukaemia HL-60 cells.*Phytochemistry*,44(2):305-10.
- Mimaki Y, Kanmoto T, Kuroda M, Sashida Y, Satomi Y, Nishino A, Nishino H. (1996) Steroidal saponins from *Hosta longipes* and their inhibitory activity on tumour promoter-induced phospholipid metabolism of HeLa cells.*Phytochemistry*,42(4):1065-70.
- Mimaki Y, Kanmoto T, Kuroda M, Sashida Y, Nishino A, Satomi Y, Nishino H. (1995) Steroidal saponins from the underground parts of *Hosta longipes* and their inhibitory activity on tumor promoter-induced phospholipid metabolism.*Chem Pharm Bull (Tokyo)*,43(7):1190-6.

シオデ

- Sashida Y, Kubo S, Mimaki Y, Nikaido T, Ohmoto T.(1992)Steroidal saponins from *Smilax riparia* and *S. china*.*Phytochemistry*,31(7):2439-43.

サンショウ

- Hur JM, Park JG, Yang KH, Park JC, Park JR, Chun SS, Choi JS, Choi JW.(2003)Effect of methanol extract of *Zanthoxylum piperitum* leaves and of its compound, protocatechuc acid, on hepatic drug metabolizing enzymes and lipid peroxidation in rats.*Biosci Biotechnol Biochem* , 67(5):945-50.
- Cho EJ, Yokozawa T, Rhyu DY, Kim SC, Shibahara N, Park JC. (2003) Study on the inhibitory effects of Korean medicinal plants and their main compounds on the 1,1-diphenyl-2-picrylhydrazyl radical. *Phytomedicine*,10(6-7):544-51.
- Cho EJ, Yokozawa T, Rhyu DY, Kim HY, Shibahara N, Park JC.(2003) The inhibitory effects of 12 medicinal plants and their component compounds on lipid peroxidation. *Am J Chin Med*,31(6):907-17.
- Jeong CH, Shim KH.(2004) Tyrosinase inhibitor isolated from the leaves of *Zanthoxylum piperitum*.*Biosci Biotechnol Biochem* , 68(9):1984-7.
- Epple G, Bryant BP, Mezine I, Lewis S. (2001)*Zanthoxylum piperitum*, an Asian spice, inhibits food intake in rats. *J Chem Ecol*., 27(8):1627-40.
- Epple G, Bryant BP, Mezine I, Lewis S.(2004) *Zanthoxylum piperitum* (DC), a potential feeding deterrent for mammals: studies with *Microtus ochrogaster* (Wagner). *Pest Manag Sci*., 60(7):624-30.
- Hashimoto K, Satoh K, Kase Y, Ishige A, Kubo M, Sasaki H, Nishikawa S, Kurosawa S, Yakabi K, Nakamura T.(2001) Modulatory effect of aliphatic acid amides from *Zanthoxylum piperitum* on isolated gastrointestinal tract. *Planta Med* ,67(2):179-81.

アケビ

Jung HJ, Lee CO, Lee KT, Choi J, Park HJ. (2004) Structure-activity relationship of oleanane disaccharides isolated from Akebia quinata versus cytotoxicity against cancer cells and NO inhibition. *Biol Pharm Bull.*, 27(5):744-7.

Mimaki Y, Kuroda M, Yokosuka A, Harada H, Fukushima M, Sashida Y. (2003) Triterpenes and triterpene saponins from the stems of Akebia trifoliata. *Chem Pharm Bull (Tokyo)*, 51(8):960-5.

ジュンサイ

石沢, 清(1963) ジュンサイ(蕪菜)の粘質物に関する研究(I) 粘質物の構成糖類、アミノ酸および灰分の検索. 山形大學紀要. 教育科學卷号 3(2), 27-36.

石沢, 清(1963) ジュンサイ(蕪菜)の粘質物に関する研究(II) 粘質物主要成分の化学組成. 山形大學紀要. 教育科學, 3(3), 89-96.

クレソン

Godz Y, Hoshino K, Akiyama H, Ishikawa T, Abe Y, Nakamura T, Otsuka H, Takeda Y, Tanimura A, Toyoda M. (1999) Constituents in watercress: inhibitors of histamine release from RBL-2H3 cells induced by antigen stimulation. *Biol Pharm Bull.* 22:1319-26.

Chiao JW, Wu H, Ramaswamy G, Conaway CC, Chung FL, Wang L, Liu D. (2004) Ingestion of an isothiocyanate metabolite from cruciferous vegetables inhibits growth of human prostate cancer cell xenografts by apoptosis and cell cycle arrest. *Carcinogenesis*, 25: 1403-8.

Hecht SS, Chung FL, Richie JP Jr, Akerkar SA, Borukhova A, Skowronski L, Carmella SG. (1995) Effects of watercress consumption on metabolism of a tobacco-specific lung carcinogen in smokers. *Cancer Epidemiol Biomarkers Prev.* 4:877-84.

Thomson B, Shaw I. A Comparison of Risk and Protective Factors for Colorectal Cancer in the Diet of New Zealand Maori and non-Maori. *Asian Pac J Cancer Prev.* 3:319-324.

モミジガサ

Shindo K, Kimura M, Iga M. (2004) Potent antioxidative activity of cacialol, a sesquiterpene contained in Cacalia delphinifolia Sieb et Zucc. *Biosci Biotechnol Biochem.* 68:1393-4.

イタドリ

Park CS, Lee YC, Kim JD, Kim HM, Kim CH. (2004) Inhibitory effects of *Polygonum cuspidatum* water extract (PCWE) and its component resveratrol [correction of rasveratrol] on acyl-coenzyme A-cholesterol acyltransferase activity for cholesterol ester synthesis in HepG2 cells. *Vascul Pharmacol.* 40(6):279-84.

Banerjee S, Bueso-Ramos C, Aggarwal BB. (2002) Suppression of 7,12-dimethylbenz(a)anthracene-induced mammary carcinogenesis in rats by resveratrol: role of nuclear factor-kappaB, cyclooxygenase 2, and matrix metalloprotease 9. *Cancer Res.* 62(17):4945-54.

Bagchi D, Das DK, Tosaki A, Bagchi M, Kothari SC. (2001) Benefits of resveratrol in women's health. *Drugs Exp Clin Res.* 27(5-6):233-48.

Lee SH, Ryu SY, Kim HB, Kim MY, Chun YJ. (2002) Induction of apoptosis by 3,4'-dimethoxy-5-hydroxystilbene in human promyeloid leukemic HL-60 cells. *Planta Med.* 68(2):123-7.

Matsuda H, Shimoda H, Morikawa T, Yoshikawa M. (2001) Phytoestrogens from the roots of *Polygonum cuspidatum* (Polygonaceae): structure-requirement of hydroxyanthraquinones for estrogenic activity. *Bioorg Med Chem Lett.* 23:11(14):1839-42.

Kimura Y, Okuda H. (2001) Resveratrol isolated from *Polygonum cuspidatum* root prevents tumor growth and metastasis to lung and tumor-induced neovascularization in Lewis lung carcinoma-bearing mice. *J Nutr.* 131(6):1844-9.

Sato M, Maulik G, Bagchi D, Das DK. (2000) Myocardial protection by protykin, a novel extract of trans-resveratrol and emodin. *Free Radic Res.* 32(2):135-44.